

Before using, please be sure to read this instruction manual. Also, store it carefully so you can read it when necessary.

第2類医薬品
鎮咳去痰薬

せき、たん、のどに

CGM クラサービッツアトローチ

Coughing is a significant physiological symptom related to the respiratory system. Continuous coughing can be distressing. Many people frequently experience symptoms such as hoarseness, throat discomfort, and throat pain due to inflammation of the throat.

Furthermore, when speaking in front of others because of phlegm, it can be very troublesome to cough continuously. This product is a pale green, aromatic, sweet ring-shaped lozenge. By slowly dissolving it in the mouth, it effectively soothes coughs, helps with phlegm, and excellently alleviates symptoms such as hoarseness, throat pain, throat swelling, throat irritation, and discomfort caused by throat inflammation.

Precautions for Use

Things you should NOT do

(If not followed, your current symptoms may worsen, and side effects or accidents are more likely to occur.)

1. Do not take if :

(1) You have previously had an allergic reaction to this product or any of its ingredients.

2. While using this product, do not use any of the following medications

Other cough suppressants and expectorants, cold medicines, sedatives, or oral medicines containing antihistamines, such as those for nasal allergies, motion sickness medicines, allergy medicines, etc.

Consultation is advised

1. Before use, the following people should consult a doctor, pharmacist, or registered seller

(1) Those under medical treatment.

(2) Pregnant women or those who believe they might be pregnant.

(3) Individuals who have experienced allergic reactions due to medications.

(4) Those with the following symptom: High fever.

2. If the following symptoms appear after use, there is a possibility of side effects. Stop using immediately and consult a doctor, pharmacist, or registered seller with this instruction manual."

Related Parts	Symptoms
Skin	Rash · Redness, Itchiness
Digestive system	Nausea · Vomiting, Loss of appetite
Nervous system	Dizziness
Respiratory system	Difficulty breathing, Shortness of breath

Rarely, the following severe symptoms may occur. In such cases, please seek immediate medical attention.

Symptom Name	Symptoms
Shock (Anaphylaxis)	Immediately after use, symptoms such as skin itching, hives, hoarse voice, sneezing, throat irritation, difficulty breathing, palpitations, and clouded consciousness may appear.

3. If the symptoms do not improve after using 5-6 times, discontinue use and consult with a doctor, pharmacist, or registered seller with this instruction manual."

Please be sure to read the reverse side as well

【Efficacy · Effect】

For the following single dose, hold it in your mouth, and let it dissolve slowly without chewing

【Dosage · Quantity】

For the following single dose, hold it in your mouth, and let it dissolve slowly without chewing

Age	Single Dose	Times per Day	Interval
Adults (15 years and older)	1 tablet	6 times	Every 2 hours or more
11 to under 15 years		4 times	
8 to under 11 years			3 times
5 to under 8 years			
Under 5 years			Do not use.

<Notes related to Dosage · Quantity>

1. Strictly adhere to the prescribed dosage and quantity.
2. When given to children, it should be under the supervision of a guardian.
3. This product is a lozenge, so do not chew or swallow it.
4. Do not give to children under 5 years of age.
5. How to take out the lozenge
Press the raised part of the PTP sheet where the lozenge is contained, break the aluminum foil on the back, and take it out. (Swallowing it as is may lead to unexpected accidents, like it getting stuck in the esophagus.)



(Diagram for removing from the PTP sheet)

【Ingredients and Action】

6 tablets (1 tablet: 1.2g) contain the following ingredients:

Ingredient	Amount	Action
Methylbisine Dium Chloride Hydrate	6mg	Acts as an antiseptic and prevents bacterial infections.
Phenol Phthalic Acid & Dextromethorphan	60mg	Suppresses the cough center and stops the cough.
Guaiacol Sulfonate Lithium	140mg	Promotes secretion in the airway, dilutes the viscosity of phlegm, making it easier to expel.

Other additives include refined sugar, powdered maltose, gelatin, Macrogol 6000, Povidone, Copper Kulkulfurinatum, Stearic Acid Calcium, Vanillin, 1-Menthyl, and fragrances.

【Storage and Handling Precautions】

1. Store in a cool, dry place away from direct sunlight
2. Keep out of reach of children.
3. Do not transfer to another container. (Mistakes in usage or changes in quality may occur.)
4. This product is hygroscopic. If the aluminum bag is opened and left for a few days, it may change color in humid conditions. Be especially careful about moisture, and do not use discolored products.
5. Do not use products past their expiration date (listed on the outer box).

For inquiries about this product, you can contact the pharmacy or drugstore where you purchased it, or the "Customer Inquiry Center" below.

Seller: **CGM Innovation Co.,Ltd.**

Address: 〒110-0003 3-9-16 Negishi, Taito-ku, Tokyo, Japan

<https://www.cgmgroup.jp>

Customer Inquiry Center: TEL: 03-6811-2345. Operating hours: 9:00-17:00 (excluding Saturdays, Sundays, and public holidays).